

Percolation in the canonical ensemble

Hao Hu¹, Henk W. J. Blöte² and Youjin Deng^{1*}

¹ *Hefei National Laboratory for Physical Sciences at Microscale,
Department of Modern Physics, University of Science
and Technology of China, Hefei 230027, China and*

² *Instituut Lorentz, Leiden University,
P.O. Box 9506, 2300 RA Leiden, The Netherlands*

(Dated: January 3, 2013)

Abstract

We study the bond percolation problem under the constraint that the total number of occupied bonds is fixed, so that the canonical ensemble applies. We show via an analytical approach that at criticality, the constraint can induce new finite-size corrections with exponent $y_{\text{can}} = 2y_t - d$ both in energy-like and magnetic quantities, where $y_t = 1/\nu$ is the thermal renormalization exponent and d is the spatial dimension. Furthermore, we find that while most of universal parameters remain unchanged, some universal amplitudes, like the excess cluster number, can be modified and become non-universal. We confirm these predictions by extensive Monte Carlo simulations of the two-dimensional percolation problem which has $y_{\text{can}} = -1/2$.

PACS numbers: 05.50.+q, 64.60.ah, 64.60.an, 64.60.F-

* yjdeng@ustc.edu.cn

I. INTRODUCTION

The bond-percolation model [1] can be considered as the $q \rightarrow 1$ limit of the q -state Potts model [2, 3]. Consider a lattice $G \equiv (V, E)$ with V (E) the vertex (edge) set, the reduced (divided by kT) Hamiltonian of the Potts model reads

$$\mathcal{H}(K, q) = -K \sum_{ij \in E} \delta_{\sigma_i \sigma_j} \quad (\sigma = 1, \dots, q), \quad (1)$$

where K is the coupling strength. By introducing bond variables on the edges and integrating out the spin degrees of freedom, one can map the Potts model onto the random-cluster (RC) model. This is the celebrated Kasteleyn-Fortuin transformation [4]. The partition sum of the RC model thus assumes the form

$$\mathcal{Z}_{\text{rc}}(u, q) = \sum_{\mathcal{A} \subseteq G} u^{|\mathcal{A}|} q^{k(\mathcal{A})} \quad (u = e^K - 1), \quad (2)$$

where the sum is over all the spanning subgraphs $\mathcal{A}$ of G , $|\mathcal{A}|$ is the number of occupied bonds in $\mathcal{A}$, and $k(\mathcal{A})$ is the number of connected components (clusters). The Kasteleyn-Fortuin mapping provides a generalization of the Potts model to non-integer values $q > 0$. In the $q \rightarrow 1$ limit, the RC model reduces to the bond percolation problem [3, 4]. In this limit, the partition sum assumes the non-singular form $(u + 1)^{|E|}$, where $|E|$ is the number of the lattice edges. Nevertheless, rich physics still exists, e.g. in the derivatives of the partition sum involving q .

Percolation in two dimensions has been extensively studied. Percolation thresholds on many two-dimensional lattices are exactly known or have been determined to very high precision (see e.g. Ref. [5] and references therein). Most of the critical exponents are also exactly known. For instance, the leading and subleading thermal renormalization exponents are $y_t = 1/\nu = 3/4$ and $y_i = -2$, respectively; the leading two critical exponents in the magnetic sector are $y_h = d - \beta/\nu = 91/48$ and $y_{h2} = 71/96$ [6, 7]. There still exist some critical exponents whose exact values are unknown, including the backbone exponent and the shortest-path fractal dimension [8, 9]. Much progress has recently been made in the context of the stochastic-Löwner evolution [10, 11].

The present work provides a study of another aspect of percolation. We introduce an “energy-like” constraint that the total number of occupied bonds is fixed and study the effects of such a constraint on the critical finite-size scaling (FSS). Since occupied bonds can

formally be considered as particles, we shall refer to percolation under this constraint as percolation in the canonical ensemble, and to the case without the constraint as percolation in the grand-canonical ensemble. Two of us have, for several years, been studying the effects of such constraints [12–14]. The leading FSS in systems under energy-like constraints is derived on the basis of the so-called Fisher renormalization [15] procedure, which was originally formulated for systems in the thermodynamic limit. Consider an energy-density-like observable ε in the critical RC model that scales as $\langle \varepsilon(L) \rangle = \varepsilon_a + \varepsilon_s L^{y_t - d}$ in the grand-canonical ensemble, where L is the linear system size and d is the spatial dimensionality. The constraint in the canonical ensemble implies:

- (i), $\varepsilon(L)$ for a given size L is fixed at the expectation value ε_a in the thermodynamic limit. This expectation value is normally different from the grand-canonical finite-size average $\langle \varepsilon(L) \rangle$.
- (ii), fluctuations of ε are forbidden—i.e., $\langle \varepsilon^2 \rangle = \langle \varepsilon \rangle^2$.

The effect of (i) is accounted for by the Fisher renormalization. The bond density $|\mathcal{A}|/|E|$ is such an observable. For bond percolation, however, since the bond variables on different edges are independent of each other, $\rho_b = \langle |\mathcal{A}| \rangle / |E|$ does not depend on the system size L —i.e., if we write $\rho_b = \rho_{b,a} + \rho_{b,s} L^{y_t - d}$, then the amplitude $\rho_{b,s} = 0$. Thus, percolation provides an ideal system to study the effects of energy-like constraints due to the suppression of fluctuations.

The remainder of this work is organized as follows. Section II describes the sampled quantities and the FSS of physical observables for percolation in the grand-canonical ensemble. Section III derives the effects of the “energy-like” constraint in the critical FSS. The numerical results are presented in Sec. IV. A brief discussion is given in Sec. V.

II. SAMPLED QUANTITIES AND FINITE-SIZE SCALING

A. Simulation and sampled quantities

We consider the bond-percolation problem on an $L \times L$ square lattice with periodic boundary conditions. The grand-canonical simulations follow the standard procedure: each edge is occupied by a bond with probability p , after which the percolation clusters are constructed. For simulations in the canonical ensemble, a Kawasaki-like scheme [16] is used.

Given an initial configuration with a total number of occupied bonds $|\mathcal{A}| = p|E|$, where $|E| = 2L^2$, an update is defined as the random selection of two edges and the exchange of their occupation states. Each sweep consists of $2L^2$ updates. Quantities are sampled after every sweep. Note that although the bond number $|\mathcal{A}| = p|E|$ happens to be an integer for the critical square-lattice bond percolation, $p|E|$ is generally a real number. In this case, one can simulate at the ceiling and the floor integer of $p|E|$ and apply an interpolation.

Given a configuration $\mathcal{A}$ as it occurs in the simulation, we denote the sizes of clusters as $\mathcal{C}_i$ ($i = 1, \dots, k(\mathcal{A})$); $\mathcal{C}_1$ is reserved for the largest cluster. The following observables were sampled.

1. *Energy-like quantities:*

- The bond-occupation density $\rho_b = \langle |\mathcal{A}| \rangle / |E|$.
- The cluster-number density $\rho_k = \langle k(\mathcal{A}) \rangle / |V|$.

2. *Specific-heat-like quantities:*

- $C_b = L^{-d} (\langle |\mathcal{A}|^2 \rangle - \langle |\mathcal{A}| \rangle^2)$.
- $C_k = L^{-d} (\langle k(\mathcal{A})^2 \rangle - \langle k(\mathcal{A}) \rangle^2)$.
- $C_{kb} = L^{-d} (\langle k(\mathcal{A}) \cdot |\mathcal{A}| \rangle - \langle k(\mathcal{A}) \rangle \langle |\mathcal{A}| \rangle)$.
- $C_2 = L^{-d} (\langle [|\mathcal{A}|/2 + k(\mathcal{A})]^2 \rangle - \langle |\mathcal{A}|/2 + k(\mathcal{A}) \rangle^2) = C_k + C_{kb} + C_b/4$.

Here, the factor $1/2$ in the definition of C_2 arises from the fact that the critical line $u_c(q)$ of the RC model (2) on the square lattice is described by $u_c(q) = q^{1/2}$, which leads to a weight $q^{|\mathcal{A}|/2 + k(\mathcal{A})}$ for a subgraph $\mathcal{A}$ along the critical line. In the canonical ensemble, the bond number $|\mathcal{A}|$ is fixed, and C_b and C_{kb} reduce to 0.

3. *Magnetic quantities:*

- The largest-cluster size $S_1 = \langle \mathcal{C}_1 \rangle$.
- The cluster-size moments S_ℓ ($\ell > 1$). Defining $\mathcal{S}_\ell = \sum_i^k \mathcal{C}_i^\ell$, we sampled $S_2 = \langle \mathcal{S}_2 \rangle$ and $\langle 3\mathcal{S}_2^2 - 2\mathcal{S}_4 \rangle$.

4. *Dimensionless quantities:*

- Wrapping probabilities R_1, R_b, R_x , and R_e . The probability R_1 counts the events that configuration $\mathcal{A}$ connects to itself along the x direction, but not along the y direction; R_b is for simultaneous wrapping in both directions; R_x is for the x direction, irrespective of the y direction; R_e is for wrapping in at least one direction. They are related as $R_e = 2R_1 + R_b$ and $R_x = R_1 + R_b$, thus only two of them are independent [17, 18].
- Universal ratio $Q_S = \langle \mathcal{C}_1^2 \rangle / S_1^2$.
- Universal ratio $Q_m = \langle \mathcal{S}_2 \rangle^2 / \langle 3\mathcal{S}_2^2 - 2\mathcal{S}_4 \rangle$.

B. Finite-size scaling in the grand-canonical ensemble

The critical FSS of the sampled quantities can be obtained from derivatives of the free-energy density $f = -L^{-d} \ln \mathcal{Z}_{\text{rc}}$ ($\mathcal{Z}_{\text{rc}}$ is given by Eq. (2)) with respect to the thermal scaling field t , the magnetic scaling field h , or the parameter q . In the grand-canonical ensemble, the FSS of $f(q, t, h, L)$ is expected to behave as

$$f(q, t, h, L) = f_r(q, t, h) + L^{-d} f_s(q, tL^{y_t}, hL^{y_h}, 1), \quad (3)$$

where higher-order scaling fields have been neglected, and f_r and f_s denote the regular and the singular part of the free-energy density, respectively. The thermal scaling field t is approximately proportional to $u - u_c$ in Eq. (2), where $u_c(q) = \sqrt{q}$ is the critical line of the q -state RC model on the square lattice.

FSS for energy-like quantities. From the partition sum in Eq. (2) and Eq. (3), it can be derived that at criticality

$$(-q)(df_c/dq) = L^{-d} \langle |\mathcal{A}|/2 + k(\mathcal{A}) \rangle \equiv \rho_b(L) + \rho_k(L) = \rho_{b,0} + \rho_{k,0} + bL^{-d}, \quad (4)$$

$$(-u/2)(\partial f/\partial u) = 2^{-1} L^{-d} \langle |\mathcal{A}| \rangle \equiv \rho_b(L) = \rho_{b,0} + aL^{y_t-d}, \quad (5)$$

$$(-q)(\partial f/\partial q) = L^{-d} \langle k(\mathcal{A}) \rangle \equiv \rho_k(L) = \rho_{k,0} - aL^{y_t-d} + bL^{-d}, \quad (6)$$

where $f_c = f(u = u_c)$ is the free-energy density along the critical line $u_c(q) = \sqrt{q}$. The last equality in Eq. (4) reflects the analyticity of f_c (including the amplitude of its finite-size dependence) along the critical line [19]. Moreover, the correction amplitude b is universal [19, 20], which is also referred to as the excess cluster number for percolation. The term $\rho_{k,0}$

follows from the exact results for the critical Potts free energy on the square lattice [21]. An exact result is also available for the triangular lattice [22]. The last equality in Eq. (5) arises from Eq. (3), and $\rho_{b,0}$ accounts for the background contribution; the self-duality of the square-lattice model yields $\rho_{b,0}(q) = 1/2$. In the $q \rightarrow 1$ limit, the amplitude a vanishes as $a \sim q - 1$. Therefore, one has for critical percolation

$$\rho_b(L) = \rho_{b,0} , \quad \rho_k(L) = \rho_{k,0} + bL^{-d} . \quad (7)$$

FSS for specific-heat-like quantities. Similarly, one has the second derivatives (at $q = 1$) as

$$- \left(q \frac{d}{dq} \right)^2 f_c \equiv C_2 = C_{2,0} + cL^{-d} , \quad (8)$$

$$- \left(u \frac{\partial}{\partial u} \right)^2 f \equiv C_b = C_{b,0} , \quad (9)$$

$$- \left(u \frac{\partial}{\partial u} \right) \left(q \frac{\partial}{\partial q} \right) f \equiv C_{kb} = C_{kb,0} + 2a' L^{y_t-d} + c' L^{-d} , \quad (10)$$

$$- \left(q \frac{\partial}{\partial q} \right)^2 f \equiv C_k = C_{k,0} - 2a' L^{y_t-d} + c'' L^{-d} , \quad (11)$$

where the last equalities in Eq. (10) and (11) arise from the fact that the parameter a in Eqs. (5) and (6) behaves as $a \approx a'(q - 1)$. The analyticity of f_c along the critical line [19] is also reflected in Eq. (8). It is also derived that the amplitude c is a universal quantity [19], called the excess fluctuation of the number $|\mathcal{A}|/2 + k(\mathcal{A})$ for critical bond percolation.

FSS for magnetic quantities. The critical FSS of magnetic quantities can be obtained by differentiating the free-energy density with respect to the magnetic scaling field h . It can be shown that at criticality,

$$S_1(L) \sim L^{y_h} , \quad S_2(L) \sim L^{2y_h} . \quad (12)$$

Corrections to FSS. At criticality, the asymptotic behavior of a quantity $O(L)$ is supposed to follow the form

$$O(L) = L^\psi (O_0 + \text{corrections}) , \quad (13)$$

where ψ is the leading critical exponent and O_0 is the amplitude. For our observables in bond percolation, values of ψ are given by the previous analysis ($\psi = 0$ for dimensionless quantities). The FSS theory predicts several types of “corrections” in Eq. (13) (see e.g. Refs.[23] and [24] for a review and references therein). These terms include corrections

from the irrelevant scaling fields; the leading one has exponent $y_i = -2$ for percolation. A regular background term, which appears e.g. as L^{-y_h} for S_1 and L^{d-2y_h} for S_2 , contributes to the “corrections”. These regular background terms are also present in the associated dimensionless ratios. For the energy-like quantities (ρ_k and ρ_b) and the specific-heat-like quantities, ψ is negative, thus the regular background (which is $L^{-\psi}$ here) included in the “corrections” in Eq. (13) is actually not a correction, instead it describes the leading behavior, which is the thermodynamic limit of the sampled quantity.

III. FINITE-SIZE SCALING IN THE CANONICAL ENSEMBLE

As mentioned in the Introduction, the critical FSS for bond percolation in the canonical ensemble is due to the suppression of the fluctuation of occupied-bond density ρ_b . In this section, we demonstrate how the critical FSS is affected by such a suppression.

Without the constraint, any configuration $\mathcal{A}$ with bond number $|\mathcal{A}| = N_b$ occurs with probability $p^{N_b}(1-p)^{(N_e-N_b)}$ ($N_e = |E|$). Thus, the value O_g of an observable O in the grand-canonical ensemble can be written as

$$\begin{aligned} O_g(p, L) &= \sum_{N_b=0}^{N_e} p^{N_b}(1-p)^{N_e-N_b} \sum_{\mathcal{A}: |\mathcal{A}|=N_b} O(\mathcal{A}) \\ &= \sum_{N_b=0}^{N_e} p^{N_b}(1-p)^{N_e-N_b} \frac{N_e!}{N_b!(N_e-N_b)!} O_c(\rho, L) , \end{aligned} \quad (14)$$

where $\rho = N_b/N_e$ is the occupied-bond density and O_c is obtained by averaging over all the configurations $\mathcal{A}$ with $|\mathcal{A}| = N_b$, of which the total number is $N_e!/N_b!(N_e-N_b)!$. Namely, O_c is the canonical-ensemble value of observable O . For a sufficiently large system, the binomial distribution in Eq. (14) is well approximated by a Gaussian distribution

$$f(p, \rho, L) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(\rho-\bar{p})^2/2\sigma^2} , \quad \sigma = \sqrt{p(1-p)/N_e} , \quad (15)$$

where $\bar{p} = p$ is the average bond density for occupation probability p , and the variance σ decreases as $\propto 1/\sqrt{N_e} \propto L^{-d/2}$. Thus, Eq. (14) can be re-written as

$$O_g(p, L) = \int d\rho f(p, \rho, L) O_c(\rho, L) . \quad (16)$$

In principle, based on the known scaling behavior of O_g in the grand-canonical ensemble, the critical FSS of O_c can be obtained by the inverse orthogonal transformation of Eq. (16). Here, we take an approximation approach by Taylor expanding $O_c(\rho, L)$ in Eq. (16) in powers of $\delta\rho = \rho - \bar{\rho}$ about $\rho = \bar{\rho} = p$, and term-by-term evaluation of the integrals:

$$\int d\rho f(p, \rho, L) O_c(\rho, L) = O_c(\bar{p}, L) + O'_c \langle \delta\rho \rangle_f + O''_c \langle (\delta\rho)^2/2 \rangle_f + \dots \quad (17)$$

where the derivatives O'_c and O''_c are taken at $\bar{p}$, and the average $\langle \rangle_f$ is over the Gaussian distribution in Eq. (15). It is easily derived that $\langle \delta\rho \rangle_f = 0$ and $\langle (\delta\rho)^2/2 \rangle_f = p(1-p)/2N_e$. Combination of Eqs. (16) and (17) yields

$$O_g(p, L) \simeq O_c(\bar{p}, L) + BO''_c L^{-d}, \quad (18)$$

with $B = p(1-p)/4$ a non-universal constant.

Near criticality p_c , we denote $\Delta p = p - p_c$. Since $\bar{p} = p$, we can equivalently denote the distance to the percolation threshold as $\Delta\bar{p} = \bar{p} - \bar{p}_c$ with $\bar{p}_c = p_c$ and $\Delta\bar{p} = \Delta p$. Thus

$$O_g(\Delta p, L) = O_{g,r}(\Delta p) + L^\psi [O_{g,s}(L^{y_t} \Delta p) + O_{g,i} L^{y_i}], \quad (19)$$

$$O_c(\Delta\bar{p}, L) = O_{c,r}(\Delta p) + L^{\psi'} [O_{c,s}(L^{y_\rho} \Delta p) + B_{\text{can}} L^{y_{\text{can}}} + O_{c,i} L^{y_i}], \quad (20)$$

where new finite-size corrections with exponent y_{can} are allowed in Eq. (20). It is known from Eq. (20) that $O''_c(\bar{p}_c, L) = O''_{c,r} + L^{\psi'+2y_\rho} O''_{c,s}$. Taylor-expanding the r.h.s of Eqs. (19-20) and substituting them in Eq. (18) give

$$O_{g,r}(0) + L^\psi [O_{g,s}(0) + O'_{g,s} L^{y_t} \Delta p + O_{g,i} L^{y_i} + \dots] = \quad (21)$$

$$O_{c,r}(0) + L^{\psi'} [O_{c,s}(0) + O'_{c,s} L^{y_\rho} \Delta p + O_{c,i} L^{y_i} + BO''_{c,s} L^{2y_\rho-d} + B_{\text{can}} L^{y_{\text{can}}} + BO'_{c,r} L^{-d-\psi'} + \dots]$$

Assuming that $2y_\rho - d < 0$, the term with $O''_{c,s}$ acts as a finite-size correction. By comparing the scaling behavior of the l.h.s and the r.h.s. of Eq. (21), we obtain

1. $\psi' = \psi$ and $y_\rho = y_t$,
2. $O_{c,r}(0) = O_{g,r}(0)$, $O_{c,s}(0) = O_{g,s}(0)$ (for $\psi \neq -d$), and $O'_{g,s} = O'_{c,s}$ etc. In other words, the scaling functions in Eqs. (19) and (20) are identical up to some correction terms.
3. new correction terms appear in the canonical ensemble, and they have exponents $2y_t - d$ and $-d - \psi$ (if $-d - \psi < 0$). This is because the leading finite-size correction

in the l.h.s of Eq. (21) is described by an exponent $y_i = -2$, and thus the terms with $L^{2y_\rho-d} = L^{2y_t-d}$ and $L^{-d-\psi'} = L^{-d-\psi}$ have to be cancelled by the newly included correction terms with $L^{y_{\text{can}}}$. It can be shown that corrections with exponent $n(2y_t-d)$ with $n = 2, 3, \dots$ can also occur if higher-order terms are kept in the above Taylor expansions.

4. for the case of $\psi = -d$, $O_{c,s}(0) = O_{g,s}(0)$ does not hold. Instead, one has $O_{g,s}(0) = O_{c,s}(0) + BO''_{c,r}$. This predicts that the excess cluster number b in Eq. (7) is changed by the constraint; further, since B and $\rho''_{c,r}$ are non-universal, the canonical-ensemble value of b is no longer universal.

Note that the assumption $2y_\rho - d < 0$ holds since $2y_t - d < 0$ for percolation in two and higher dimensions.

Finally, we mention that although the constraint is “energy-like”, the derivation applies to both the energy-like and the magnetic observables.

IV. NUMERICAL RESULTS

To examine the above theoretical derivations, we carry out Monte Carlo simulations for the critical bond percolation on the square lattice, with a fixed bond number $N_b = p_c|E| = L^2$, and for 21 sizes in range $4 \leq L \leq 4000$. The number of samples was about 10^8 for $L \leq 480$, 5×10^7 for $L = 800$, 2.5×10^7 for $L = 1600$, and 10^6 for $L = 4000$. For the purpose of comparison, additional simulations were also carried out in the grand-canonical ensemble.

A. Evidence for the correction exponent $y_{\text{can}} = 2y_t - d$

We first examine the critical FSS of the wrapping probabilities. In the grand-canonical ensemble, the finite-size corrections arise only from the irrelevant scaling fields for which the leading exponent $y_i = -2$. This is illustrated in the inset of Fig. 1. In the canonical ensemble, the existence of the newly induced correction exponent $y_{\text{can}} = 2y_t - d = -1/2$ is clearly demonstrated by the approximately linear behavior for large L in Fig. 1, where R_1 is plotted versus $L^{-1/2}$.

According to the least-squares criterion, we fitted the data for these wrapping probabilities




FIG. 1: (Color online). Plot of the wrapping probability R_1 versus $L^{-1/2}$ in the canonical ensemble (main plot). The inset shows R_1 versus L^{-2} in the grand-canonical ensemble. The straight lines represent the leading finite-size dependence as obtained from the fit. The error bars of the data are smaller than the size of the data points.

by

$$O = O_0 + B_1 L^{y_{\text{can}}} + B_2 L^{2y_{\text{can}}} + B_i L^{y_i}, \quad (22)$$

where O_0 represents the universal value for $L \rightarrow \infty$, and the correction exponent y_i is fixed at -2 . As a precaution against higher-order correction terms not included in the fit formula, the data points for small $L < L_{\text{min}}$ were gradually excluded to see how the residual χ^2 changes with respect to L_{min} . In general, we use results of the fit corresponding to a L_{min} for which the quality of fit is reasonable, and for which subsequent increases of L_{min} do not cause the χ^2 value to drop by vastly more than one unit per degree of freedom. In practice, the word “reasonable” means here that χ^2 is less than or close to the number of degrees of freedom, and that the fitted parameters become stable.

The results are shown in Table I. The error margins are quoted as two times of the statistical errors in the fits, which also applies to other tables, in order to account for possible systematic errors. The universal values O_0 can be exactly obtained [25, 26]. In the grand-canonical ensemble, these exact values of O_0 were used and the amplitudes B_1 and B_2 were set at 0, so that the fit formula in Eq. (22) has only a single free parameter. Indeed, such a simple formula can well describe the data with $L \geq 16$ for all the wrapping

TABLE I: Fit results for wrapping probabilities. The exact values of O_0 [25, 26], and $y_i = -2$ are used here. The abbreviations “GE” and “CE” stand for grand- and canonical ensemble, respectively. Entries “-” indicate the absence of a result, and the numbers without error bars are fixed in the fits. Error margins are quoted as two times of the statistical errors in the fits.

Fit using $O = O_0 + B_1 L^{y_{\text{can}}} + B_2 L^{2y_{\text{can}}} + B_i L^{y_i}$								
		O_0	B_1	y_{can}	B_2	B_i	y_i	L_{min}
R_1	GE	0.169 415 435	—	—	—	0.23(1)	−2	16
		0.169 415 435	—	—	—	0.24(3)	−2.02(4)	8
	CE	0.169 5(4)	0.11(2)	−0.53(5)	—	—	—	160
		0.169 415 435	0.091 (8)	−0.502 (13)	0.08(2)	0.56(7)	−2	12
		0.169 415 435	0.089 2(5)	−1/2	0.082 (4)	0.55(3)	−2	12
R_b	GE	0.351 642 855	—	—	—	−0.254 (11)	−2	12
		0.351 642 855	—	—	—	−0.30(5)	−2.06(8)	8
	CE	0.351 5(6)	−0.10(3)	−0.55(7)	—	—	—	120
		0.351 642 855	−0.075 (11)	−0.50(2)	−0.07(2)	−0.53(5)	−2	8
		0.351 642 855	−0.075 8(8)	−1/2	−0.066 (5)	−0.53(3)	−2	8
R_x	GE	0.521 058 290	—	—	—	−0.021 (16)	−2	16
		0.521 058 290	—	—	—	−0.1(1)	−2.4(6)	8
	CE	0.521 1(2)	0.020 (4)	−0.56(8)	—	—	—	32
		0.521 058 290	0.018 (7)	−0.54(8)	—	—	—	120
		0.521 058 290	0.014 (9)	−0.51(9)	0.016(13)	—	—	16
		0.521 058 290	0.016 (12)	−0.52(11)	0.01(3)	0.03(8)	−2	10
		0.521 058 290	0.013 4(8)	−1/2	0.016 (6)	0.02(4)	−2	10
R_e	GE	0.690 473 725	—	—	—	0.22(2)	−2	16
		0.690 473 725	—	—	—	0.18 (4)	−2.0(1)	8
	CE	0.690 5(6)	0.13(3)	−0.53(6)	—	—	—	120
		0.690 473 725	0.13(3)	−0.53(3)	—	0(7)	−2	120
		0.690 473 725	0.10(3)	−0.50(3)	0.13(4)	—	—	40
		0.690 473 725	0.12(3)	−0.52(3)	0.05(9)	0.9(5)	−2	20
		0.690 473 725	0.102(2)	−1/2	0.10(2)	0.5(3)	−2	24




FIG. 2: (Color online). Plot of $\Delta R_x = R_x - O_0 - B_1 L^{y_{\text{can}}}$ versus $L^{2y_{\text{can}}}$ for the canonical wrapping probability R_x . The straight line represents the leading finite-size dependence of ΔR_x , i.e. $B_2 L^{2y_{\text{can}}}$. Parameter O_0 and y_{can} are fixed at the exact value and $-1/2$ respectively, and the amplitudes B_1 , B_2 are obtained from the fit.

probabilities (R_1 , R_b , R_x , and R_e); further, for R_x the amplitude of B_i is very small. As expected, including terms with B_1 and/or B_2 only yields messy information and does not improve the quality of the fits. In the canonical ensemble, we also fitted the data by Eq. (22) with a single correction term $B_1 L^{y_{\text{can}}}$ ($B_2 = B_i = 0$). As shown in Tab. I, the data up to sufficiently large L_{min} (except R_x) have to be discarded for a reasonable residual χ^2 . Nevertheless, we find that (i), the correction term has an exponent close to $-1/2$, and (ii), the estimates of O_0 agree well with the exact values. A better description of the R data can be obtained by including correction terms with B_i and/or B_2 ; for simplicity the exact values of O_0 are used. As a result, the estimates of y_{can} become more accurate, and are in good agreement with the prediction $y_{\text{can}} = -1/2$. We note that the correction coefficient B_i takes different values in the grand-canonical and the canonical ensemble. This is also in agreement with the theoretical expectation. As predicted in Sec. III, in addition to those with exponent $n(2y_t - d)$ ($n = 1, 2, \dots$), correction terms with exponent $-d - \psi$ can also exist in the canonical ensemble. For the wrapping probabilities, one has $\psi = 0$ and thus such a correction term has the same exponent as $B_i L^{y_i}$. Finally, in order to illustrate the term with $L^{2y_{\text{can}}} (= L^{-1})$, we plot $\Delta R_x = R_x - O_0 - B_1 L^{y_{\text{can}}}$ versus $L^{2y_{\text{can}}}$ as in Fig. 2.

TABLE II: Fit results for S_1 , S_2 , and the universal ratios Q_S and Q_m .

Fit using $O = L^\psi(O_0 + B_1 L^{y_{\text{can}}} + B_r L^{y_r} + B_i L^{y_i})$							
		O_0	B_1	y_{can}	B_r	B_i	L_{min}
S_1	GE	0.984 48(3)	—	—	1.7(13)	−2(2)	32
	CE	0.984 3(4)	0.066 (14)	−0.50(5)	6(5)	−7(6)	24
S_2	GE	1.035 97(4)	—	—	1.7(7)	−3.0(14)	32
	CE	1.035 8(5)	0.05(3)	−0.51(12)	5(3)	−7(5)	24
Q_S	GE	0.960 17(1)	—	—	1.4(8)	−1.8(12)	40
	CE	0.960 0(3)	0.072 (8)	−0.48(4)	—	—	200
		0.960 03(13)	0.075 (5)	−0.49(2)	−0.3(16)	0(2)	24
Q_m	GE	0.870 56(2)	—	—	−1.1(3)	1.92(6)	32
	CE	0.870 4(2)	0.138 (7)	−0.491 (13)	−1.1(6)	1.3(9)	16

In order to demonstrate that the correction exponent for the grand-canonical wrapping probabilities is indeed $y_i = -2$, we also fitted the data by Eq. (22) with y_i being free, O_0 being fixed at the exact values and $B_1 = B_2 = 0$. The fit results are also included in Table I. The result is $-2.02(4)$, $-2.06(8)$, $-2.4(6)$ and $-2.0(1)$, for R_1 , R_b , R_x and R_e , respectively. These results are in compatible with the existing numerical result $y_i = -2.003(5)$ [18] for R_1 , and agree well with the exact value $y_i = -2$. It is interesting to mention that although correction of form $L^{-2}(1 + A \log L)$ (A is a constant) has been observed for other observables in the square-lattice bond percolation [24], there is no evidence for such a logarithmic factor in the finite-size corrections for the wrapping probabilities.

For magnetic quantities, we consider the largest-cluster size S_1 and the second cluster-size moment S_2 , as well as the dimensionless ratios. We fitted the data by

$$O = L^\psi(O_0 + B_1 L^{y_{\text{can}}} + B_r L^{y_r} + B_i L^{y_i}), \quad (23)$$

with $y_i = -2$ and ψ being fixed at the exactly known value for the two-dimensional percolation. The term with L^{y_r} is the correction from the regular background, with $y_r = -y_h = -91/48$ for S_1 and Q_S , and $y_r = d - 2y_h = -43/24$ for S_2 and Q_m . The results are shown in Table II. Again, we find that (i), the values of O_0 remain unchanged

TABLE III: Fit results for the cluster-number density ρ_k as determined for the bond (BP) and site (SP) percolation problem. Some fits make use of the exact results $\rho_{k,0}^{\text{bond}} = 0.098\,076\,211$ [21] and $b_g = 0.883\,576\,308$ [19].

Fit using $\rho_k = \rho_{k,0} + L^{-2}(b + B_1 L^{y_{\text{can}}} + B_i L^{y_i})$							
		$\rho_{k,0}$	b	B_1	y_{can}	B_i	L_{min}
BP	GE	0.098 076 22(7)	0.883 6(6)	—	—	0.18(3)	6
		0.098 076 211	0.883 576 308	—	—	0.178 (14)	6
	CE	0.098 076 21(4)	0.23(2)	0.57(5)	−0.47(6)	0.0(4)	8
		0.098 076 211	0.226 (8)	0.564 (12)	−0.47(3)	—	10
		0.098 076 211	0.233 (4)	0.60(2)	−1/2	—	24
SP	GE	0.027 598 03(2)	0.883 5(2)	—	—	−0.17(4)	16
		0.027 598 02(2)	0.883 576 308	—	—	−0.22(3)	20
	CE	0.027 598 00(4)	0.41(2)	0.64(7)	−0.46(7)	—	24
		0.027 597 99(4)	0.417 (2)	0.687 (11)	−1/2	—	24

in the canonical ensemble, irrespective of whether they are universal (for Q_S and Q_m) or non-universal (for S_1 and S_2), and (ii), new correction terms with an exponent $y_{\text{can}} = -1/2$ are introduced. The estimate $Q_{m,0} = 0.870\,56(2)$ in the grand-canonical ensemble agrees with the existing result $0.870\,53(2)$ [14].

B. Change of the universal excess cluster number

For energy-like quantities, we consider the cluster-number density ρ_k , and analyze the data by

$$\rho_k = \rho_{k,0} + L^{-2}(b + B_1 L^{y_{\text{can}}} + B_i L^{y_i}), \quad (24)$$

with $y_i = -2$ being fixed. The background term can be exactly obtained as $\rho_{k,0}^{\text{bond}} = (3\sqrt{3} - 5)/2$ for the critical square-lattice bond percolation [21]. In the grand-canonical ensemble, the excess cluster number b is known to be universal and the value has been obtained as $b_g = 0.883\,576\,308$ [19], with subscript g for the grand-canonical ensemble.

The fit results are given in Table III. As in the above subsection, the correction term




FIG. 3: (Color online). Plot of the cluster-number density ρ_k versus L^{-2} for the bond percolation model. The straight lines represent the leading finite-size dependence of ρ_k as obtained from the fit.

with $y_{\text{can}} = -1/2$ is observed in the canonical ensemble, and the background contribution $\rho_{k,0}$ remains unchanged. However, the excess cluster number is now $b_c = 0.233(4)$, clearly different from the universal value $b_g = 0.883\,576\,308$ [19] in the grand-canonical ensemble. This indeed confirms the theoretical prediction for the case of $\psi = -2$ in Sec. III. In other words, one expects that $b_g = b_c + B\rho''_{k,r}$, where $B = p(1-p)/4$ is a constant and $\rho''_{k,r}$ is the second derivative $\rho''_{k,r}$ of the regular part of ρ_k with respect to the bond density. Since both B and $\rho''_{k,r}$ are non-universal, the canonical-ensemble value b_c is *no longer* universal. As an illustration, we plot ρ_k versus L^{-2} in Fig. 3, where the difference of b_g and b_c is reflected by the different slopes of the data lines.

In order to examine the “*non-universal*” nature of the excess cluster number b_c in the canonical ensemble, we also performed simulations of square-lattice site-percolation problem at the percolation threshold $p_c^{\text{site}} = 0.592\,746\,02$ [5, 24, 27]. In the grand-canonical ensemble, the simulations used 15 system sizes in range $4 \leq L \leq 512$, and the results of the fits by Eq. (24) are given in Tab. III. The estimate of $b_g = 0.883\,5(2)$ agrees well with the universal value $0.883\,576\,308$ [19].

In the canonical ensemble, the total number of occupied sites $N_s(L) = p_c^{\text{site}} L^2$ is fixed. However, $N_s(L)$ is not an integer, and thus the actual simulations were carried out for the total occupied-site number $[N_s(L)]$ and $[N_s(L)] + 1$, with $[]$ for the floor integer. The

Monte Carlo results at $N_s(L)$ were then obtained by linear interpolation. The simulations used 16 system sizes L in range $4 \leq L \leq 1024$. The results of fits using Eq. (24) are also given in Tab. III. As expected, one has a correction term with exponent $-1/2$ and $b_c^{(\text{site})} [= 0.417(2)] \neq b_g$. The “*non-universal*” property of b_c is demonstrated by the fact $b_c^{(\text{site})} \neq b_c^{(\text{bond})}$.

In addition, from our fits, one gets $\rho_{k,0}^{\text{site}} = 0.027\,598\,00(5)$, which is in good agreement with the existing result $0.027\,598\,1(3)$ [20], and reduces the error margin significantly.

We also study the FSS of the specific-heat-like quantity C_2 for the bond percolation model and find that due to the limited precision, the data for $L \geq 8$ are well described by $C_2(L) = C_{2,0} + cL^{-2}$. As already mentioned in Section II, the excess fluctuation c is universal. In the grand-canonical ensemble, the fit yields $C_{2,0} = 0.039\,446\,(4)$ and $c_g = 0.105\,5(7)$; the former is consistent with the existing result $C_{2,0} = 0.039\,44(4)$ [28], the latter agrees well with the exactly known value $c = 0.105\,436\,634$ [19]. In the canonical ensemble, the result is $C_{2,0} = 0.039\,446\,(6)$ and $c_c = -0.33(5)$. It is interesting to observe that not only the magnitude of the excess fluctuation c is modified, but that also its sign has changed.

C. Universality of scaling functions

In Sec. III, we state that the scaling functions in Eqs. (19) and (20) are identical up to some correction terms, which reflects the universality of the functions. In order to demonstrate this, we carried out simulations near the critical point. We plot the grand-canonical R_x versus $L^{y_t}(p - p_c)$ and the canonical R_x versus $L^{y_t}(\bar{p} - \bar{p}_c)$ as in Fig. 4. As expected, data points in both the ensembles nicely collapse to the same curve.

V. DISCUSSION

We derive the critical finite-size scaling behavior of percolation under the constraint that the total number of occupied bonds/sites is fixed, and confirm theoretical predictions for the two-dimensional percolation by means of Monte Carlo simulation in two dimensions. In particular, it is found that with the constraint, new finite-size corrections with exponent $n(2y_t - d)$ ($n = 1, 2, \dots$) are induced and the excess cluster number becomes non-universal. We note that our theory in Sec. III can be used to explain the observed correction expo-




FIG. 4: (Color online). Plot of R_x versus t for bond percolation. Parameter t represents $L^{y_t}(p - p_c)$ in the grand-canonical ensemble (GE), and $L^{y_t}(\bar{p} - \bar{p}_c)$ in the canonical ensemble (CE). The excellent collapse demonstrates the universality of scaling functions near the critical point.

nent ≈ -0.5 for the canonical wrapping probabilities in simulating the two-dimensional percolation by the Newman-Ziff algorithm [17, 18]. The predictions should be valid in any dimension with $d \geq 2$. We believe that this work provides an additional useful reference for percolation—a pedagogical system in the field of statistical mechanics. Furthermore, our work can help to understand the critical finite-size-scaling properties of other statistical systems in the canonical ensemble with a fixed total number of particles, which is the usual situation during experiments.

Acknowledgments

This work was supported by NSFC (under grant numbers 10975127, 91024026 and 11275185), CAS, and Fundamental Research Funds for the Central Universities No. 2340000034. We would like to thank R. M. Ziff for helpful comments. H. Hu is grateful for the hospitality of the Instituut Lorentz of Leiden University.

[1] Stauffer D and Aharony A 1994 *Introduction to Percolation Theory* revised 2nd edn (London: Taylor & Francis)

- [2] Potts R B 1952 *Proc. Cambridge Philos. Soc.* **48** 106
- [3] Wu F Y 1978 *J. Stat. Phys* **18** 115
- [4] Kasteleyn P W and Fortuin C M 1969 *J. Phys. Soc. Japan* **26** (Suppl.) 11
Fortuin C M and Kasteleyn P W 1972 *Physica (Amsterdam)* **57** 536
- [5] Ziff R M 2011 *Phys. Procedia* **15** 106
- [6] Nienhuis B 1987 *Phase Transitions and Critical Phenomena* vol 11 ed Domb C and Lebowitz J L (London: Academic) p 1
- [7] Cardy J L 1987 *Phase Transitions and Critical Phenomena* vol 11 ed Domb C and Lebowitz J L (London: Academic) p 55
- [8] Grassberger P 1992 *J. Phys. A* **25** 5475
Grassberger P 1992 *J. Phys. A* **25** 5867
Grassberger P 1999 *Physica A* **262** 251
- [9] Zhou Z, Yang J, Deng Y and Ziff R M 2012 *arXiv:1112.3428* and references therein
- [10] Kager W and Nienhuis B 2004 *J. Stat. Phys.* **115** 1149
- [11] Cardy J L 2005 *Ann. Phys.* **318** 81
- [12] Deng Y and Blöte H W J 2004 *Phys. Rev. E* **70** 046111
- [13] Deng Y, Heringa J R and Blöte H W J 2005 *Phys. Rev. E* **71** 036115
- [14] Deng Y and Blöte H W J 2005 *arXiv:cond-mat/0508348*
- [15] Fisher M E 1968 *Phys. Rev.* **176** 257
- [16] Kawasaki K 1972 *Phase Transitions and Critical Phenomena* vol 2 ed Domb C and Green M (London: Academic) p 443
- [17] Newman M E J and Ziff R M 2000 *Phys. Rev. Lett.* **85** 4104
- [18] Newman M E J and Ziff R M 2001 *Phys. Rev. E* **64** 016706
- [19] Kleban P and Ziff R M 1998 *Phys. Rev. B* **57** R8075
- [20] Ziff R M, Finch S R and Adamchik V S 1997 *Phys. Rev. Lett.* **79** 3447
- [21] Temperley H N V and Lieb E H 1971 *Proc. R. Soc. Lond. A.* **322** 251
- [22] Baxter R J, Temperley H N V and Ashley S E 1978 *Proc. R. Soc. Lond. A.* **358** 535
- [23] Garoni T M, Ossola G, Polin M and Sokal A D 2011 *J. Stat. Phys.* **144** 459
- [24] Feng X, Deng Y and Blöte 2008 *Phys. Rev. E* **78** 031136
- [25] Pinson H T 1994 *J. Stat. Phys.* **75** 1167
- [26] Ziff R M, Lorenz C D and Kleban P 1999 *Physica A* **266** 17-26

- [27] Lee M J 2008 *Phys. Rev. E* **78** 031131
- [28] Deng Y and Yang X 2006 *Phys. Rev. E* **73** 066116